

Short communications

Acoustic concomitants of the defensive discharges of a primitive bombardier beetle (*Metrius contractus*)*

T. Eisner¹, J. Yack² and D. J. Aneshansley³

¹Department of Neurobiology and Behavior, W347 Mudd Hall, Cornell University, Ithaca, NY 14853, USA

²Department of Biology, College of Natural Sciences, Carleton University, Ottawa, Ontario, Canada K1S 5B6

³Department of Agricultural and Biological Engineering, Riley-Robb Hall, Cornell University, Ithaca, NY 14853, USA

Summary. *Metrius contractus*, a primitive paussoid bombardier beetle, emits its defensive quinonoid froth with accompanying sound (a faint “hiss”), but the sound is not pulsed, indicating that the secretory emission itself is not pulsed. Pulsed secretory delivery in bombardier beetles appears to occur in Brachinini only.

Key words. Defensive glands – discharge mechanism – quinones – bioacoustics – Carabidae

Arthropods that discharge defensive secretions usually do so inaudibly. The exceptions include a number of insects that discharge froths. The grasshopper, *Romalea guttata*, for instance, emits a bubbling foam when disturbed, a mixture of secretion and tracheal air that it ejects with a hissing sound from the mesothoracic spiracles (Fig. 1A) (Alsop 1970). Defensive froth emission also occurs in many arctiid moths, which discharge the foam from the anterolateral corners of the thorax (Fig. 1B), and may do so with accompanying sound (Carpenter 1938). But sound producers also include insects that eject jets of fluid rather than froth. Loudest, perhaps, are the familiar bombardier beetles of the genus *Brachinus*, which emit a distinct “pop” when they discharge their hot quinonoid spray (Schildknecht 1957; Aneshansley *et al.* 1969). Naturalists early on called attention to the detonations. In what is probably one of the first accounts of *Brachinus*, a Swedish student, Daniel Rolander, reports encountering a remarkable insect, which upon being seized, gave off a “kleinen Knalle” (a little crack) and a puff of “Rauch” (smoke). He called the insect a “Schussfliege” (or shooting fly; the term bombardiers was not coined until later), told Linnaeus about it, and published a description of it in 1750, in the Proceedings of the Royal Swedish Academy of Sciences [available to us in the German translation (Rolander 1754)].

Bombardier beetles comprise two evolutionary branches (see Fig. 1 in Eisner *et al.* 2001) within the family Carabidae (references in Eisner *et al.* 2001). One branch, the brachinoid branch, includes the “advanced” bombardiers of the tribe Barachinini (comprising the genera *Brachinus* and *Stenaptinus*, among others) and the primitive species of the tribe Crepidogastrini. The other branch, the paussoid branch, features a number of tribes judged to be more or less advanced (Mystropomini, Ozaenini, Paussini), and one tribe, the Metriini, generally considered to be the most primitive of the paussoids. There is uncertainty whether the two branches are linked at their base (in other words, whether the two lineages have a common bombarding ancestor) or whether they represent parallel lineages in which the ability to bombard evolved independently (references in Eisner *et al.* 2001).

To help settle the question it becomes useful to learn as much as possible about the two groups that are the most primitive within each branch, the Crepidogastrini and the Metriini. Beetles of both these tribes are relatively hard to come by, but thanks to helpful collectors we have been able to obtain live individuals of each, and have published descriptions of their defenses (Eisner *et al.* 2000, 2001).

One finding of interest concerned the evolution of an important feature of the discharge mechanism, namely the pulsed nature of the secretory expulsion. The Brachinini typically show such pulsed emission (Fig. 8 in Eisner *et al.* 2001), which represents an adaptive refinement of the delivery system (Dean *et al.* 1990). The crepidogastrines, we have now shown, do not pulse their emissions (Eisner *et al.* 2001), indicating that the ability to pulse did not evolve in brachinoids until *after* the appearance of the brachinine stock. The absence of pulsation in crepidogastrines was detected acoustically, from the absence of temporal discontinuity in the sound oscillograms of these beetles.

When we published on *Metrius* (Eisner *et al.* 2000), we had not recorded its discharge sounds, and were therefore unable to determine whether this beetle pulses its discharges. Given the primitive phyletic status of *Metrius*, and the fact that in other paussoids (Ozaenini) the discharge sounds failed to reveal pulsation (T.

Correspondence to: T. Eisner, e-mail: te14@cornell.edu

* Paper No. 182 in the series “Defense mechanisms of Arthropods”. Paper No. 181 is Eisner T, Aneshansley DJ, Yack J, Attygalle AB, Eisner M, Spray mechanism of crepidogastrine bombardier beetles (Carabidae: Crepidogastrini). Chemoecology 11:209–219




Fig. 1 **A.** *Romalea guttata*, discharging defensive froth (arrow) from the mesothoracic spiracle. **B.** The arctiid moth, *Utetheisa ornatrix*, discharging froth from the cervical region. Many arctiids emit such froth with accompanying sound. *U. ornatrix* does so inaudibly. **C, D.** Tethered *Metrius contractus*, in rear end view, discharging from the same gland on two separate occasions. In **(C)** the photo caught the emission early on, after onset of the discharge, and the froth has just formed; in **(D)** the picture is of the discharge at a later stage, when many tiny droplets, engendered by the bursting bubbles of the froth, are spewed into the air. Bars **(A)** 5 mm; **(B, C)** 2 mm

Eisner & D. J. Aneshansley, unpublished recordings of *Goniotropis nicaraguensis*), we thought it unlikely that *Metrius* would pulse. However, without evidence we could not be certain. We have now made recordings with *Metrius* and here report that its discharges, like that of crepidogastrines and probably paussoids generally, is non-pulsed. We are thus led to the conclusion that pulsation evolved in the brachinoid branch only, and that within the brachinoids it is an exclusive feature of the Brachiniini.

Our sound recordings were done with *Metrius contractus*, using the same apparatus and techniques we used in recording the discharge sounds of crepidogastrines (Eisner *et al.* 2001). We had four live *M. contractus* available and made sound recordings with each. It was clear just from listening to the discharge sounds of *Metrius* that these were more faint and of different quality than the ones typical of *Brachinus*. The *Metrius* sound, unlike the sharp “pop” emitted by *Brachinus*, is in the nature of a hiss. The difference is doubtless a consequence of the different ways in which the two beetles effect their discharges. While *Brachinus* emits its secretion as discrete forcible jets (Eisner and Aneshansley 1999), *Metrius* discharges a froth that bubbles away on the surface of its body (Eisner *et al.* 2000) (Fig. 1C, D). To the ear, in fact, a *Metrius* discharge has very much the attributes of a faint *Romalea* emission.

We recorded several discharges from each of the four *M. contractus*. The beetles were tethered as de-

scribed for other bombardiers (Eisner *et al.* 2001) and they were caused to discharge by pinching their legs with forceps. The periods over which they were subjected to stimulation ranged from 1.5 to 3 minutes. The interval between consecutive discharges was therefore in the range of seconds to over a minute. *M. contractus* discharges from one gland at a time, restricting its emission to the gland of the body side stimulated (Eisner *et al.* 2000). Since we usually pinched the legs of a beetle in no particular order, we had no way of




Fig. 2 Oscillograms of sound accompanying discharges from a single *Metrius contractus*. **A, B,** and **C** represent the first, third, and fifth discharges from a single gland over a period of 40 seconds. The relative sound amplitudes decrease, while the durations increase, with successive discharges. Bar 50 ms

determining the sequence of alternation of the discharges in the oscillograms or whether any one discharge stemmed from the right or left gland. It was impossible therefore to determine in most cases how the acoustical properties of the discharges of any one gland changed as a function of the progressive depletion of that gland, although we could tell that toward the end of the sound tracts, when both glands must have been approaching depletion, the discharges tended to become longer and of decreased intensity, as if the glands were then "fizzing out." This was clearly evident from the oscillogram of the one beetle in which we did record 5 consecutive discharges from one gland. Three of the discharge sounds of that beetle are depicted in Fig. 2. At any rate, the tracts left no doubt as regards pulsation, for which there was no evidence in any of the 13 discharges that we recorded.

A question that has never been answered is whether the accompanying sound contributes to the defensive effectiveness of a bombardier's ejections. While one can easily imagine the relatively loud detonations of a *Brachinus* having a startling effect on a predator, one is hard put envisioning the soft hiss of a *Metrius* having an impact on any but the smallest of enemies.

Acknowledgements

The study was supported by grant AI02908 from the

Received 8 July 2001; accepted 23 July 2001.

National Institutes of Health. We are deeply indebted to Kipling Will for sending us the *M. contractus* from California.

References

- Alsop DW (1970) Defensive glands of arthropods: comparative morphology of selected types. Doctoral thesis, Cornell University
- Aneshansley DJ, Eisner T, Widom JM, Widom B (1969) Biochemistry at 100°C: explosive secretory discharge of bombardier beetles (*Brachinus*). *Science* 165:61–63
- Carpenter H (1938) Audible emission of defensive froth by insects. *Proc Zool Soc* 108:243–252
- Dean J, Aneshansley DJ, Edgerton HE, Eisner T (1990) Defensive spray of the bombardier beetle: a biological pulse jet. *Science* 248:1219–1221
- Eisner T, Aneshansley DJ (1999) Spray aiming in the bombardier beetle: photographic evidence. *Proc Natl Acad Sci USA* 96:9705–9709
- Eisner T, Aneshansley DJ, Eisner M, Attygalle AB, Alsop DW, Meinwald J (2000) Spray mechanism of the most primitive bombardier beetle (*Metrius contractus*). *J Exp Biol* 203:1265–1275
- Eisner T, Aneshansley DJ, Yack J, Attygalle AB, Eisner M (2001) Spray mechanism of crepidogastrine bombardier beetles (Carabidae; Crepidogastrine). *Chemoeology* 11:209–219
- Rolander D (1754) Die Schussfliege. *Abhand Konigl Schwed Akad Wissensch* 12:298–302
- Schildknecht H (1957) Zur Chemie des Bombardierkäfers. *Angew Chem* 69:62